


Oncogene Surveillance: Differential role of Runx3 and p53

Yoshiaki Ito, MD PhD

Professor, Cancer Science Institute of Singapore



RUNX family are evolutionarily conserved metazoan genes that encode a DNA binding subunit of a transcription factor. Recently, RUNX gene along with p53 and other transcription factors were found to be present in unicellular organism considered as a precursor to the metazoan. RUNX is known to function in cell specification in development and frequently involved in carcinogenesis. We reported earlier that Runx3 functions as a barrier to gastric and colon carcinogenesis (1,2). More recently, we began analyzing the K-Ras-induced lung cancer using mouse model.

In an attempt to cure cancer by restoration of p53, two groups performed sophisticated mouse studies using K-Ras-induced lung adenocarcinoma. The results reported in 2010 indicated that p53 destroyed only high-grade adenocarcinoma but not adenoma (3,4). If p53 is entirely responsible for oncogene surveillance, the results described in these two papers are puzzling.

We used mice carrying Runx3^{fl/fl}, K-Ras^{LSL-G12D}, Adeno-Cre or tm/CreERT in different combinations and conditionally activated Cre recombinase by two different methods: one is to express in a large number of lung cells, the other in much smaller number of cells. We found that K-Ras^{G12D} alone induced lung adenocarcinoma when Cre was activated in a large number of cells.

However, If Cre was activated in a small number of cells, activation of K-Ras^{G12D} alone or inactivation of Runx3 alone did not induce any tumor over one year. Only when Cre was activated in a small number of cells in the mouse carrying K-Ras^{G12D}, Runx3^{fl/fl}, lung adenocarcinoma developed very rapidly, indicating that inactivation of Runx3 is required for K-Ras to induce adenocarcinoma.

We analyzed the possible mechanism for this phenomenon and the results will be presented. This study suggested that first line of defense against tumor formation is carried out by Runx3 and the second line by p53.

Acknowledgement: This study was performed in close collaboration with my former colleague, Dr. Suk-Chul Bae in Chungbuk National Univ, Korea.

References:

- 1) Ito et al, Gastroenterology 140:1536-1546, 2011
- 2) Ito et al, Cancer Cell 14:226-237, 2008
- 3) Junttila et al, Nature 468:567-571, 2010
- 4) Feldser et al, Nature 468:572-575, 2010

Yoshiaki Ito

Senior Principal Investigator, Cancer Science Institute Singapore
National University of Singapore
Yong Loo Lin Professor of Medical Oncology
Department of Medicine, Yong Loo Lin School of Medicine,
National University of Singapore
Email: csiitoy@nus.edu.sg

POSITIONS AND HONORS

1968	Kuroya Award (Japanese Society for Microbiology)
1995	Princess Takamatsu Cancer Research Award
2003	Tomizo Yoshida Prize (Japanese Cancer Association)
2010	President's Science Awards (2010) --- (A*STAR, Singapore)

RECENT PUBLICATIONS

1. Voon DC, Wang H, Koo Jk, Nguyen TA, Hor YT, Chu YS, Ito K, Fukamachi H, Chan SL, Thierry JP, Ito Y. Runx3 protects gastric epithelial cells against epithelial-mesenchymal transition-induced cellular plasticity and tumorigenicity. *Stem Cells*, 2012 Oct;30 (10):2088-99
2. Linda Shyue Huey Chuang , Soak Kuan Lai , Maki Murata-Hori , Ayumi Yamada , Hoi-Yeung Li , Jayantha Gunaratne, Ito Y (2012) RUNX3 interactome reveals novel centrosomal targeting of RUNX family of transcription factors. *Cell Cycle*, 15 May 2012;11(10):1938-1947
3. Ito K, Chuang LS, Ito T, Chang TL, Fukamachi H, Salto-Tellez M, Ito Y. Loss of Runx3 is a key event in inducing precancerous state of the stomach. *Gastroenterology*. (2011) Jan;140:1536-1546
4. Chang TL, Ito K, Ko TK, Liu Q, Yang S, Salto-Tellez M, Yeoh KG, Fukamachi H, Ito Y. Claudin-1 has tumor suppressive activity and is a direct target of RUNX3 in gastric epithelial cells. *Gastroenterology*, (2010) Jan;138(1):255-65
5. Ito K, Lim AC, Salto-Tellez M, Motoda L, Osato M, Chuang LS, Lee CW, Voon DC, Koo JK, Wang H, Fukamachi H, Ito Y. *RUNX3 Attenuates beta-Catenin/T Cell Factors in Intestinal Tumorigenesis. Cancer Cell*. (2008) Sep 9; 14(3): 226-37.
6. Ito, K., Liu, Q., Salto-Tellez, M., Yano, T., Tada, K., Ida, H., Huang, C., Shah, N., Inoue, M., Rajnakova, A., Hiong, K.C, Peh, B.K, Han, H.C, Ito, T., Teh, M., Yeoh, K.G. and Ito, Y. (2005) RUNX3, a novel tumor suppressor, is highly inactivated in gastric cancer by protein mislocalization. *Cancer Res* 65, 7743-7750
7. Li, Q.L., Ito, K., Sakakura, C., Fukumachi, H., Inoue, K., Chi, X.Z., Lee, K.Y., Nomura, S., Lee, C.W., Han, S.B., Kim, H.M., Kim, W.J., Yamamoto, H., Yamashita, N., Yano, T., Ikeda, T., Itohara, S., Inazawa, J., Abe, T., Hagiwara, A., Yamagishi, H., Ooe, A., Kaneda, A., Sugimura, T., Ushijima, T., Bae, S.C., Ito, Y. (2002). Casual Relationship between the Loss of RUNX3 Expression and Gastric Cancer. *Cell* 109, 113-124.
8. Inoue, K., Ozaki, S., Shiga, T., Ito, K., Masuda, T., Okado, N., Iseda, T., Kawaguchi, S., Ogawa, M., Bae, S.C., Yamashita, N., Itohara, S., Kudo, N., Ito, Y. (2002). RUNX 3 controls the axonal projection of proprioceptive dorsal root ganglion neurons. *Nature Neurosci* 5, 946-954
9. Ogawa, E., Maruyama, M., Kagoshima, H., Inuzuka, M., Lu, J., Satake, M., Shigesada, K., and Ito, Y. (1993). PEBP2/PEA2 represents a family of transcription factors homologous to the products of the *Drosophila runt* gene and the human AML1 gene. *Proc. Natl. Acad. Sci. USA* 90, 6859-6863.
10. Smart, J. E., and Ito, Y. (1978). Three species of polyoma virus tumor antigens share common peptides probably near the amino termini of the proteins. *Cell* 15, 1427-1437.